
Table 4. Estimated numbers of diagnoses of AIDS in children <13 years of age, by year of diagnosis and exposure category, 1998–2002—United States

	Year of diagnosis					Cumulative
	1998	1999	2000	2001	2002	through 2002
Race/ethnicity						
White, not Hispanic	33	19	13	16	7	1,606
Black, not Hispanic	156	129	85	70	58	5,476
Hispanic	47	33	18	22	24	2,111
Asian/Pacific Islander	0	1	2	2	1	59
American Indian/Alaska Native	1	0	0	0	0	31
Exposure category						
Hemophilia/coagulation disorder	0	0	0	0	0	236
Mother with, or at risk for, HIV infection	236	181	115	106	90	8,629
Injection drug use	46	43	20	13	9	3,309
Sex with injection drug user	31	24	15	5	5	1,526
Sex with bisexual male	6	6	2	3	4	202
Sex with person with hemophilia	1	2	0	1	0	35
Sex with HIV-infected transfusion recipient	1	0	0	0	0	26
Sex with HIV-infected person, risk not specified	62	50	36	32	32	1,453
Receipt of blood transfusion, blood components, or tissue	0	1	2	2	3	159
Has HIV infection, risk not specified	89	55	40	50	37	1,920
Receipt of blood transfusion, blood components, or tissue	1	0	1	0	1	390
Other/risk not reported or identified	0	2	2	3	0	45
Total ^b	238	183	118	110	92	9,300

Note. These numbers do not represent actual cases in persons with a diagnosis of AIDS. Rather, these numbers are point estimates of cases diagnosed that have been adjusted for reporting delays and for redistribution of cases in persons initially reported without an identified risk. The estimates have not been adjusted for incomplete reporting.

Figure 1. Estimated numbers of diagnoses of AIDS in children <13 years of age, by year of diagnosis, 1992–2002—United States



Note. These numbers do not represent actual cases in children with a diagnosis of AIDS. Rather, these numbers are point estimates of cases diagnosed that have been adjusted for reporting delays. The estimates have not been adjusted for incomplete reporting. The 1993 AIDS surveillance case definition change affected only cases in adults and adolescents, not the cases in children.

^a Includes children with a diagnosis of AIDS, from the beginning of the epidemic through 2002.

^b Includes children of unknown or multiple race. Cumulative total includes 17 children of unknown or multiple race. Because column totals were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.